

§ 201.56–12

- (1) Cotyledons:
 - (i) Less than half of the original cotyledon tissue remaining attached.
 - (ii) Less than half of the original cotyledon tissue free of necrosis or decay.
- (2) Epicotyl:
 - (i) Missing. (May be assumed to be present if cotyledons are intact.)
 - (ii) [Reserved]
- (3) Hypocotyl:
 - (i) Deep open cracks or grainy lesions extending into the conducting tissue.
 - (ii) Malformed, such as markedly shortened, curled, or thickened.
 - (iii) Watery.
- (4) Root:
 - (i) None.
 - (ii) Weak, stubby, or missing primary root with weak secondary or adventitious roots.
- (5) Seedling:
 - (i) One or more essential structures impaired as a result of decay from primary infection.
 - (ii) Albino.

[59 FR 64506, Dec. 14, 1994]

§ 201.56–12 Miscellaneous plant families.

Kinds of seed by family:

- Carrot family, Apiaceae (Umbelliferae)—carrot, celery, celeriac, dill, parsley, parsnip;
 - Hemp family, Cannabaceae—hemp;
 - Dichondra family, Dichondraceae—dichondra;
 - Geranium family, Geraniaceae—alfilaria;
 - Mint family, Lamiaceae (Labiatae)—sage, summer savory; benne family, Pedaliaceae—sesame;
 - Rose family, Rosaceae—little burnet;
 - Nightshade family, Solanaceae—eggplant, tomato, husk tomato, pepper, tobacco; and
 - Valerian family, Valerianaceae—cornsalad.
- (a) General description.
 - (1) Germination habit: Epigeal dicot.
 - (2) Food reserves: Cotyledons; endosperm may or may not be present, depending on the kind.
 - (3) Shoot system: The hypocotyl elongates, carrying the cotyledons above the soil surface. The epicotyl usually does not show any development within the test period.
 - (4) Root system: A primary root; secondary roots may or may not develop

7 CFR Ch. I (1–1–05 Edition)

within the test period, depending on the kind.

- (b) Abnormal seedling description.
 - (1) Cotyledons:
 - (i) Less than half of the original cotyledon tissue remaining attached.
 - (ii) Less than half of the original cotyledon tissue free of necrosis or decay.
 - (2) Epicotyl:
 - (i) Missing. (May be assumed to be present if the cotyledons are intact.)
 - (ii) [Reserved]
 - (3) Hypocotyl:
 - (i) Malformed, such as markedly shortened, curled, or thickened.
 - (ii) Deep open cracks extending into the conducting tissue.
 - (iii) Watery.
 - (4) Root:
 - (i) None.
 - (ii) Missing or stubby primary root with weak secondary or adventitious roots.
 - (5) Seedling:
 - (i) One or more essential structures impaired as a result of decay from primary infection.
 - (ii) Albino.

[59 FR 64506, Dec. 14, 1994]

§ 201.57 Hard seeds.

Seeds which remain hard at the end of the prescribed test because they have not absorbed water, due to an impermeable seed coat, are to be counted as “hard seed.” If at the end of the germination period provided for legumes, okra, cotton and dichondra in these rules and regulations there are still present swollen seeds or seeds of these kinds which have just started to germinate, all seeds or seedlings except the above-stated shall be removed and the test continued for 5 additional days and the normal seedlings included in the percentage of germination. For flatpea, continue the swollen seed in test for 14 days when germinating at 15–25 °C or for 10 days when germinating at 20 °C.

[5 FR 33, Jan. 4, 1940, as amended at 10 FR 9952, Aug. 11, 1945; 20 FR 7936, Oct. 21, 1955; 65 FR 1708, Jan. 11, 2000]

§ 201.57a Dormant seeds.

Dormant seeds are viable seeds, other than hard seeds, which fail to germinate when provided the specified